

## June 27, 2026 CLEAR ALIGNERS: SELECTING THE PERFECT CASES

	1.	The four cases never to treat with clear alignorms. Class II, growing Class III, bilateral crossbite open bite cases.	•	g	Т	F
2.		Distalizing molars with clear aligners is a simple and predictable orthodontic movement.		Т	F	
	<ol> <li>In general, crowding, spacing, overjet, overbite and open bite can in the 2-3mm range are ideal treatment candidates for clear align therapy in a GP office.</li> <li>Clear Aligners are designed to correct skeletal issues such as Class II and Class III malocclusions.</li> </ol>					F
				as	Т	F
	<ol> <li>Utilizing functional appliances to correct Class II and Class III malocclusions is done prior to using Clear Aligners.</li> </ol>		III	Т	F	
TO RECEIVE 1 HOUR OF CONTINUING EDUCATION, PLEASE COMPLETE THE TEST SCAN AND EMAIL TO LEE@RONDEAUSEMINARS.COM.						
^iName:				ַטט	DMD	OTHER
*Address:						
*City:		*State/Prov:	*Zip/Postal: _			
*Email:		(CE will		ill be e	mailed)	
Phone:		AGD #:			_(if ap	plicable)

\*required fields

 $\square$  Yes, I would like to receive further information from Rondeau Seminars.